NASA/TM-2000-209891, Vol. 171



Technical Report Series on the Boreal Ecosystem-Atmosphere Study (BOREAS)

Forrest G. Hall and Andrea Papagno, Editors

Volume 171 BOREAS TE-17 Production Efficiency Model Images

S.J. Goetz, S.N. Goward, S.D. Prince, K. Czajkowski, and R.O. Dubayah

National Aeronautics and Space Administration

Goddard Space Flight Center Greenbelt, Maryland 20771

The NASA STI Program Office ... in Profile

Since its founding, NASA has been dedicated to the advancement of aeronautics and space science. The NASA Scientific and Technical Information (STI) Program Office plays a key part in helping NASA maintain this important role.

The NASA STI Program Office is operated by Langley Research Center, the lead center for NASA's scientific and technical information. The NASA STI Program Office provides access to the NASA STI Database, the largest collection of aeronautical and space science STI in the world. The Program Office is also NASA's institutional mechanism for disseminating the results of its research and development activities. These results are published by NASA in the NASA STI Report Series, which includes the following report types:

- TECHNICAL PUBLICATION. Reports of completed research or a major significant phase of research that present the results of NASA programs and include extensive data or theoretical analysis. Includes compilations of significant scientific and technical data and information deemed to be of continuing reference value. NASA's counterpart of peer-reviewed formal professional papers but has less stringent limitations on manuscript length and extent of graphic presentations.
- TECHNICAL MEMORANDUM. Scientific and technical findings that are preliminary or of specialized interest, e.g., quick release reports, working papers, and bibliographies that contain minimal annotation. Does not contain extensive analysis.
- CONTRACTOR REPORT. Scientific and technical findings by NASA-sponsored contractors and grantees.

- CONFERENCE PUBLICATION. Collected papers from scientific and technical conferences, symposia, seminars, or other meetings sponsored or cosponsored by NASA.
- SPECIAL PUBLICATION. Scientific, technical, or historical information from NASA programs, projects, and mission, often concerned with subjects having substantial public interest.
- TECHNICAL TRANSLATION.
 English-language translations of foreign scientific and technical material pertinent to NASA's mission.

Specialized services that complement the STI Program Office's diverse offerings include creating custom thesauri, building customized databases, organizing and publishing research results . . . even providing videos.

For more information about the NASA STI Program Office, see the following:

- Access the NASA STI Program Home Page at http://www.sti.nasa.gov/STI-homepage.html
- E-mail your question via the Internet to help@sti.nasa.gov
- Fax your question to the NASA Access Help Desk at (301) 621-0134
- Telephone the NASA Access Help Desk at (301) 621-0390
- Write to:
 NASA Access Help Desk
 NASA Center for AeroSpace Information
 7121 Standard Drive

Hanover, MD 21076-1320

NASA/TM-2000-209891, Vol. 171



Technical Report Series on the Boreal Ecosystem-Atmosphere Study (BOREAS)

Forrest G. Hall and Andrea Papagno, Editors

Volume 171 BOREAS TE-17 Production Efficiency Model Images

Scott J. Goetz, Samual N. Goward, Stephen D. Prince, Kevin Czajkowski, and Ralph O. Dubayah University of Maryland, College Park

National Aeronautics and Space Administration

Goddard Space Flight Center Greenbelt, Maryland 20771

	Available from:	
NIAGA Conton for A G T C		Ni-diamater distribution of the control of the cont
NASA Center for AeroSpace Information		National Technical Information Service
7121 Standard Drive		5285 Port Royal Road
Hanover, MD 21076-1320		Springfield, VA 22161
Drice Code: A17		Del - C 1 - A 10
Price Code: A17		Price Code: A10

BOREAS TE-17 Production Efficiency Model Images

Scott J. Goetz, Samual N. Goward, Stephen D. Prince, Kevin Czajkowski, Ralph Dubayah

Summary

A BOREAS version of the Global Production Efficiency Model (http://www.inform.umd.edu/glopem/) was developed by TE-17 to generate maps of gross and net primary production, autotrophic respiration, and light use efficiency for the BOREAS region. This document provides basic information on the model and how the maps were generated. The data generated by the model are stored in binary image-format files.

Note that the files of this data set on the BOREAS CD-ROMs have been compressed using the Gzip program. See Section 8.2 for details.

Table of Contents

- 1) Data Set Overview
- 2) Investigator(s)
- 3) Theory of Measurements
- 4) Equipment
- 5) Data Acquisition Methods
- 6) Observations
- 7) Data Description
- 8) Data Organization
- 9) Data Manipulations
- 10) Errors
- 11) Notes
- 12) Application of the Data Set
- 13) Future Modifications and Plans
- 14) Software
- 15) Data Access
- 16) Output Products and Availability
- 17) References
- 18) Glossary of Terms
- 19) List of Acronyms
- 20) Document Information

1. Data Set Overview

1.1 Data Set Identification

BOREAS TE-17 Production Efficiency Model Images

1.2 Data Set Introduction

The Boreal Forest Production Efficiency Model (Boreal-PEM) is composed of a suite of models that provide estimates of the variables needed to drive a production efficiency model (i.e., one based on restrictions in the conversion "efficiency" of absorbed photosynthetically active radiation (APAR) in terms of unstressed gross primary production (GPP) through short-term environmental physiology).

1.3 Objective/Purpose

The purpose of the production efficiency modeling for the BOReal Ecosystem-Atmosphere Study (BOREAS) was to use remotely sensed observations to estimate GPP and net primary production (NPP) at the spatial resolution of the Advanced Very High Resolution Radiometer (AVHRR) Local Area Coverage (LAC) for the entire BOREAS region. Higher resolution maps would be possible with, e.g., Earth Observing System (EOS) instruments (launched 1998) or with sensors on aircraft platforms. The advantage of satellite data was that, once the component models were validated, consistent measurements could be made across the entire region. Thus, this approach captured gradients of land use intensity and climate. In addition to NPP and GPP, Boreal-PEM provided the data needed to model above-ground biomass and canopy conductance (e.g., maps of air temperature, vapor pressure deficit (VPD), etc.). In combination with land cover and deforestation maps, the remotely sensed measurements of NPP can yield estimates of the impact of land cover change on carbon storage, which is a focus of the follow-on work.

1.4 Summary of Parameters

A number of surface variables required to implement Boreal-PEM are retrieved on a daily basis using surface "parameter retrieval" algorithms. These include surface radiometric temperature (Ts), ambient air temperature (Ta), atmospheric precipitable water vapor amount (U), surface absolute humidity, VPD, fractional PAR absorption (FPAR), APAR, standing above-ground biomass, and a cumulative surface wetness index (CSI). Other variables related to the efficiency of light utilization, or the carbon yield of APAR, include the proportion of vegetation cover types that utilize the C3 or C4 photosynthetic pathways. This is derived using long-term climatological information and modeled biomass.

1.5 Discussion

Boreal-PEM consists of linked models of canopy radiative transfer, canopy utilization of APAR, and physical environmental variables that have a multiplicative effect on stomatal control. The model is entirely driven with satellite-retrieved surface variables (e.g., APAR, air temperature, soil moisture, absolute humidity, etc.). The resulting "stressed" GPP is reduced to NPP through carbon expenditures associated with autotrophic respiration derived from standing above-ground biomass. The model results closely approximated surface measurements of both physical and biological variables, including NPP, within the BOREAS study areas and were clearly associated with land cover type (i.e., broadleaf deciduous, needleleaf evergreen, etc.).

1.6 Related Data Sets

BOREAS Level-3b AVHRR-LAC Imagery: Scaled At-sensor Radiance in LGSOWG Format

BOREAS Level-4b AVHRR-LAC Ten-Day Composite Images: At-sensor Radiance

BOREAS Level-4c AVHRR-LAC Ten-Day Composite Images: Surface Parameters

BOREAS RSS-04 1994 Southern Study Area Jack Pine LAI and FPAR Data

BOREAS RSS-07 Regional LAI and FPAR Images From Ten-Day AVHRR-LAC Composites

BOREAS RSS-14 Level-1 GOES-7 Visible, IR and Water-vapor Images

BOREAS RSS-14 Level-1a GOES-7 Visible, IR, and Water-vapor Images

BOREAS RSS-14 Level-2 GOES-7 Shortwave and Longwave Radiation Images

BOREAS RSS-14 Level-1 GOES-8 Visible, IR and Water-vapor Images

BOREAS RSS-14 Level-1a GOES-8 Visible, IR and Water-vapor Images

2. Investigator(s)

2.1 Investigator(s) Name and Title

Samual N. Goward (PI), Professor and Chair Stephen D. Prince, Professor Scott J. Goetz, Research Scientist Kevin Czajkowski, Research Scientist Ralph O. Dubayah, Assoc. Professor

2.2 Title of Investigation

Biospheric Dynamics in the Boreal Forest Ecotone

2.3 Contact Information

Contact 1:

Dr. Scott J. Goetz Dept. of Geography University of Maryland College Park, MD 20742-8225 (301) 405-1297 (301) 314-9299 (fax) sgoetz@geog.umd.edu.

Contact 2:

Samual N. Goward Dept. of Geography University of Maryland College Park, MD 20742-8225 sg21@umail.umd.edu

Contact 3:

Stephen D. Prince Dept. of Geography University of Maryland College Park, MD 20742-8225 sp43@umail.umd.edu

Contact 4:

Kevin Czajkowski Dept. of Geography University of Maryland College Park, MD 20742-8225 kczajkow@geog.umd.edu

Contact 5:

Ralph O. Dubayah Dept. of Geography University of Maryland College Park, MD 20742-8225 rdubayah@geog.umd.edu

Contact 6:

Andrea Papagno Raytheon ITSS NASA GSFC Code 923 Greenbelt, MD 20771 (301) 286-3134 (301) 286-0239 (fax) Andrea.Papagno@gsfc.nasa.gov

3. Theory of Measurements

The model is driven primarily by vegetation light absorption, which determines potential photosynthetic rates. The potential rates are reduced by stress terms, including Ta, VPD, and CSI, which act on stomatal physiology (i.e., conductance, Gs). The "stressed" photosynthetic rates are used to provide an actual value of daily carbon gain. The loss of carbon via autotrophic respiration (Ra) is modeled based on a semi-empirical relationship with standing above-ground biomass, which, in turn, is derived from monthly minimum visible reflectance. A contextural approach is used to derive pixel-by-pixel maps of the environmental variables on a daily basis for all days when AVHRR scenes were available (the BOREAS results were based on 35 acquisitions throughout the 1994 growing season). GPP, NPP, Ra, and the amount of NPP per unit APAR on an annual basis are also generated as output maps.

Variations of the PEM approach have been taken by various investigators, beginning with correlative models first described by Kumar and Monteith (1982), Tucker et al. (1983), and Asrar et al. (1985). The Carnegie Ames Stanford Approach (CASA) (Potter et al., 1993) was the first model to use the PEM concept on a global scale, and Global PEM (GLO-PEM) was the first PEM to utilize variables retrieved entirely with remotely sensed observations on a global scale. Descriptions of individual model components and their performance have been reported in numerous journal publications (see Section 10.2.1).

4. Equipment

4.1 Sensor/Instrument Description

See associated data set documentation in Section 1.6.

4.1.1 Collection Environment

See associated data set documentation in Section 1.6.

4.1.2 Source/Platform

See associated data set documentation in Section 1.6.

4.1.3 Source/Platform Mission Objectives

See associated data set documentation in Section 1.6.

4.1.4 Key Variables

See associated data set documentation in Section 1.6.

4.1.5 Principles of Operation

See associated data set documentation in Section 1.6.

4.1.6 Sensor/Instrument Measurement Geometry

See associated data set documentation in Section 1.6.

4.1.7 Manufacturer of Sensor/Instrument

See associated data set documentation in Section 1.6.

4.2 Calibration

4.2.1 Specifications

See associated data set documentation in Section 1.6.

4.2.1.1 Tolerance

See associated data set documentation in Section 1.6.

4.2.2 Frequency of Calibration

See associated data set documentation in Section 1.6.

4.2.3 Other Calibration Information

See associated data set documentation in Section 1.6.

5. Data Acquisition Methods

All data are recovered with algorithms driven by optical and thermal AVHRR observations, with the exception of incident PAR, which is derived from the Geostationary Operational Environmental Satellite (GOES) observations (provided by Eric Smith for BOREAS).

6. Observations

6.1 Data Notes

The AVHRR data as provided by the Canada Centre for Remote Sensing (CCRS) to BOREAS were used for these analyses. No additional screening was necessary; however, Boreal-PEM utilizes a series of routines to test for the presence of subpixel clouds, and this technique was improved using the BOREAS data set (Czajkowski et al., 1997a).

6.2 Field Notes

None given.

7. Data Description

7.1 Spatial Characteristics

7.1.1 Spatial Coverage

The nominal spatial resolution of the data was 1.1 km at nadir. All data were resampled to 1-km resolution using image mapping techniques developed at CCRS (called GEOCOMP) and validated at the University of Maryland (Czajkowski et al., 1997b).

The corners of the total area covered by the model are defined by the following:

BOREAS Grid		NAD83 Coord	dinates
X (KM)	Y (KM)	Longitude	Latitude
175.000	0.000	108.51193W	50.97184N
175.000	1000.000	107.87384W	59.94373N
975.000	0.000	97.31119W	50.13370N
975.000	1000.000	93.92214W	58.89897N

The total area covered by the model runs was 800 km by 1,000 km, corresponding to the BOREAS study region within the BOREAS Information System (BORIS) X grid range of 175-975 km and the BORIS Y grid range of 0-1000 km, encompassing both the Northern Study Area (NSA) and the Southern Study Area (SSA).

Image coordinates (upper left origin) are easily related to BORIS grid coordinates (lower left origin):

```
grid\_X = image\_pixel

grid\_Y = 1000 - image\_line
```

For example, the NSA old jack pine tower site (T7Q8T) with BORIS grid coordinates Y=617, X=769 has image coordinates of line=383, pixel=769.

7.1.2 Spatial Coverage Map

Not available.

7.1.3 Spatial Resolution

All data were resampled by CCRS to 1-km pixels before being used as model input fields.

7.1.4 Projection

The area mapped is projected in the BOREAS grid projection, which is based on the ellipsoidal version of the Albers Equal-Area Conic (AEAC) projection. The projection has the following parameters:

Datum: NAD83

Ellipsoid: GRS80 or WGS84
Origin: 111.000°W 51.000°N
Standard Parallels: 52° 30' 00"N
58° 30' 00"N

Units of Measure: kilometers

7.1.5 Grid Description

Lines and pixels increase from the upper left corner of the image.

7.2 Temporal Characteristics

7.2.1 Temporal Coverage

The 34 ÅVHRR acquisitions defined the observational period during the growing season. The earliest image was 16-Apr-1994 and the latest was 07-Sep-1994. GPP, Ra, and NPP for periods between acquisitions were interpolated to a daily interval using linear interpolation. Annual (growing season) results are summed daily values.

7.2.2 Temporal Coverage Map

Not applicable.

7.2.3 Temporal Resolution

The model operates on a daily time-step. But the data represent annual values.

7.3 Input Data Characteristics

Input data required by the model are summarized in the following table.

Table 1. Boreal-PEM input variables

7.3.1 Input Parameter/ Variable	7.3.2 Variable Description/ Definition	7.3.3 Unit of Measurement	7.3.4 Data Source	7.3.5 Data Range
visible	spectral exoatmospheric reflectance	%	AVHRR channel 1	0, 25
near infrared	spectral exoatmospheric reflectance	%	AVHRR channel 2	0, 50
T4	thermal emission (brightness temperature)	degrees C	AVHRR channel 4	0, 50°
T5	thermal emission (brightness temperature)	degrees C	AVHRR channel 5	0, 50°
NDVI	normalised difference vegetation index	unitless	(ch2-ch1) / (ch2+ch1)	0, 1
mean Ta	climatological mean air temperature	degrees C	Leemans and Cramer	0, 50°
incident PAR	incident photosynthetically active radiation	MJ/day	GOES (Eric Smith/BORIS)	0, 12
e	surface spectral emissivity	unitless	Prabhakara and Dalu	0, 1

7.4 Output Data Characteristics

Model output is image format, binary single byte-per-pixel (i.e., 8-bit) values that range between 0-255. See Section 8.2 for data format, scaling, and coordinate information. The following table describes the output variables.

Table 2. Boreal-PEM output variables

7.4.1 Output Parameter/ Variable	7.4.2 Variable Description /Definition	7.4.3 Unit of Measurement	7.4.4 Data Source	7.4.5 Data Range
Ts	surface radiometric temperature	degrees C	modeled	0, 50°
Та	ambient air temperature	degrees C	modeled	0, 50°
VPD	vapor pressure deficit	millibars (mb)	modeled	0, 50
CSI	cumulative surface wetness index	unitless	modeled	-5, 5
APAR	absorbed photosynthetically active radiation	megajoules (MJ)	modeled	0, 12 / day 100, 1100 / yr

W	standing above-ground biomass	kg/m ²	modeled	0, 40
GPP*	gross primary production	gC/m ²	modeled	0, 40 / day 0, 1900 / yr
Ra*	autotrophic respiration	gC/m ²	modeled	0, 35 / day 0, 1100 / yr
NPP*	net primary production	gC/m ²	modeled	0, 20 / day 0, 850 / yr
e*	carbon yield of APAR	gC/MJ	modeled	0, 1.25

^{*} annual images provided to BORIS

7.5 Sample Data Record

Not applicable to image data.

8. Data Organization

8.1 Data Granularity

The smallest unit of data tracked by BORIS was the entire set of images.

8.2 Data Format(s)

8.2.1 Uncompressed Data Files

The entire data set contains one American Standard Code for Information Interchange (ASCII) header file and six image files. Each image file contains 1,000 records (image lines) that contain 800 1-byte pixel values.

File Num.	Description	Record Size	# Records	Bytes/Pixel
1	ASCII header file	 80	 11	N/A
2	APAR Image	800	1000	1
3	Biomass Image	800	1000	1
4	Gross Primary Production	800	1000	1
5	Light Use Efficiency	800	1000	1
6	Net Primary Production	800	1000	1
7	Autotrophic Respiration	800	1000	1

The image files values must be multiplied or divided by a scaling factor to obtain physical units. The appropriate factors are:

- apar multiply image values by 4 for actual range of 0 900 gC/m²-yr
- biomass multiply image values by 6 for actual range of 0 40 kg/m²
- gpp multiply image values by 6 for actual range of 0 1500 gC/m²-yr
- lue divide image values by 255 for actual range of 0 1 gC/MJ
- npp multiply image values by 3 for actual range of 0 600 gC/m²-yr
- resp multiply image values by 4 for actual range of 0 900 gC/m²-yr

8.2.2 Compressed CD-ROM Files

On the BOREAS CD-ROMs, files 2 through 7 listed in Section 8.2.1 have been compressed with the Gzip compression program (file name *.gz). These data have been compressed using gzip version 1.2.4 and the high compression (-9) option (Copyright (C) 1992-1993 Jean-loup Gailly). Gzip (GNU zip) uses the Lempel-Ziv algorithm (Welch, 1994) used in the zip and PKZIP programs. The compressed files may be uncompressed using gzip (-d option) or gunzip. Gzip is available from many Web sites (for example, ftp site prep.ai.mit.edu/pub/gnu/gzip-*.*) for a variety of operating systems in both executable and source code form. Versions of the decompression software for various systems are included on the CD-ROMs.

9. Data Manipulations

9.1 Formulae

See Section 9.1.1.

9.1.1 Derivation Techniques and Algorithms

$$T_s = 1.274 + \left(\frac{e''}{2}\right)T_4 + \left(\frac{e''}{2}\right)T_5$$
 (after Becker and Li, 1990)

where: T_4 = apparent temperature in AVHRR channel 4 (K)

 T_5 = apparent temperature in AVHRR channel 5 (K)

e = surface spectral emissivity;

$$e = \left(\frac{1 - e^{+}}{0.6}\right) * ND VI + \left(e^{+} - \left[\frac{0.05(1 - e^{+})}{0.6}\right]\right)$$
(2)

e* = 'grey-body' emissivity for unvegetated surface (Prabhakara et al., 1977). and e', e" are terms to characterize between-band differences in e.

$$Ta = a * 0.7 + b$$
 (3)

where: a = slope of Ts/NDVI relationship

b = intercept of Ts/NDVI relationship

Ta = intercept

where NDVI = 0.7 a,b change with moving window contextural linear regression (TVX) approach.

VPD = vapor pressure deficit (D);

$$D = 0.611* \left[\exp\left(17.27 * \frac{T_a}{T_a + 237}\right) - \exp\left(17.27 * \frac{T_d}{T_d + 237}\right) \right]$$
(4)

where T_d = dewpoint temperature;

$$T_{d} = \frac{\ln(\lambda + 1) + \ln(U) - 0.1133}{0.0393}$$
 (5)

and I = coefficient to adjust for latitude and season (Smith 1966).

U = atmospheric precipitable water content (cm);

$$U = 17.32 * \frac{(DT - 0.6831)}{(TS - 291.97)} + 0.5456$$
 (6)

and DT = T4-T5

CSI = cumulative surface wetness index (Sg);

$$\Sigma y = \Sigma y_{t-1} + y_t \tag{7}$$

where g_t = slope of Ts/NDVI at time (t) (i.e., a simple 'bucket' model in which g_t , corrected for solar zenith angle effects, varies relative to a normalized value, 0.5);

APAR = absorbed photosynthetically active radiation,

$$APAR = FPAR * PAR$$
 (8)

where: FPAR = 1.67*NDVI - 0.08

NDVI = (ch2-ch1)/(ch2+ch1)

PAR = incident PAR from GOES (E. Smith/BORIS)

NPP = net primary production;

$$NPP = GPP - Ra$$
 (9)

GPP = gross primary production;

$$GPP = APAR * e_g$$
 (10a)

where $e_g = e_g^*$ (modified by multipliers that simulate stomatal physiology, Gs); i.e., Ta, VPD, CSI)

$$\varepsilon_{\rm g}^* = 55.2\alpha$$
 (10b)

x = quantum yield of photosynthesis (Collatz, et al. 1991)

 e_n = carbon yield of APAR (g/MJ);

$$e_n = NPP / APAR \tag{11}$$

Ra = autotrophic respiration;

$$Ra = \left[\left(\frac{0.4}{0.75} \right) * \left(\frac{10^3}{10^3 W + 50} \right) \right]$$
 (12)

W = standing aboveground biomass;

$$W = 7166.1 (\rho^{-2.6})$$
 13)


r = minimized visible reflectance (ch1) on an annual basis, cloud-screened and corrected for sun angle

9.2 Data Processing Sequence

9.2.1 Processing Steps

Each individual AVHRR image is input to the model, and the various component models derive intermediate products that are then interpolated to a daily basis and integrated annually. The full processing sequence is complex because of the availability of individual AVHRR acquisitions, the number of individual component algorithms, and the exchange of variables between the different model time-steps. A sense of the processing steps can be best acquired in Prince and Goward (1995), Goetz et al. (1998), and the flowchart below.

BORIS staff copied the ASCII and compressed the binary files for release on CD-ROM.



9.2.2 Processing Changes

None given.

9.3 Calculations

9.3.1 Special Corrections/Adjustments

None.

9.3.2 Calculated Variables

See Table 2.

9.4 Graphs and Plots

For BOREAS-specific results see Goetz and Prince (1998).

10. Errors

10.1 Sources of Error

There are several potential sources of error that can affect the model results. These include errors in the data that drive the model (e.g., calibration and correction of the AVHRR reflectances and temperatures, the GOES PAR maps and their time-integration), errors in the recovery of surface variables within model component algorithms (e.g., Ts, Ta, U, VPD, FPAR, APAR, W, en, etc.), and multiplicative or canceling errors in variables derived from other recovered variables and parameters (e.g., APAR, Ra, GPP, NPP, etc.).

10.2 Quality Assessment

10.2.1 Model Validation by Source

Numerous journal articles describe efforts to test, compare, and validate various component model results. The validity of physical environmental components of the model (i.e., Ts, Ta, U, VPD, CSI) have been assessed with different field experiment data (SNF, FIFE, HAPEX-Sahel, OTTER, GEWEX, BOREAS) as listed in the following table (see acronym list, Section 19).

Author(s)	<u>Variables</u>	Field Data
Czajkowski et al. (1997a)	Ta	BOREAS
Goetz (1997)	CSI	FIFE
Goetz et al. (1998)	CSI	BOREAS
Goward et al. (1994)	Ta, U, VPD, CSI	OTTER
Goward and Dye (1997)	Ta, U, VPD, CSI	Global
Prihodko and Goward (1997)	Ta	FIFE
Prince and Goward (1995)	Ta, U, VPD, CSI	Global
Prince et al. (1998)	Ts, Ta, U, VPD	FIFE, HAPEX, GEWEX, BOREAS

The biological components of the model have been assessed in the following journal publications:

Validation of biological components of Boreal-PEM.

Author(s)	Variables	Field Data
Goetz and Prince (1996, 1998)	e _n , e _g , e _g *, NPP, GPP, Ra, APAR	SNF
Goetz et al. (1998)	e _n , NPP, GPP, Ra	BOREAS
Goward et al. (1994)	FPAR, APAR	OTTER
Hanan et al. (1995, 1997)	e _n , g _s , NPP, Ra, FPAR, APAR	HAPEX-Sahel
Prince and Goward (1995)	e _n , e _g , NPP, GPP, Ra, FPAR, APAR	Global
Prince et al. (1995)	e _n , e _g , NPP, Ra, APAR	SNF, Global

Numerous other articles have examined more general aspects of model components (particularly with respect to Ts and U). A brief summary of results from the previous tables is provided below in Sections 10.2.2-10.2.4.

10.2.2 Confidence Level/Accuracy Judgment

Confidence in the results is very high in terms of the spatial patterns and magnitudes within the images and moderate to high in terms of the absolute values of variables recovered (see Sections 10.2.3 and 10.2.4). Comparisons showed a close correspondence between measured and inferred soil moisture at the BOREAS sites. There was also good agreement between inferred and site measurements of biomass and NPP, although the biomass values were underestimated compared to those derived with an independent technique (i.e., Hall et al., 1995).

10.2.3 Measurement Error for Variables

In quantitative terms, the results of model component validation work showed that Ts could be retrieved with root mean square (RMS) errors of 3.5 °C for a range of 48 °C; Ta with 3.9 °C over a range of 36 °C; U with 0.6 cm over a range of 3.6 cm; and VPD with 10.9 mb over a range of 58 mb (Prince et al., 1998). There was some evidence of compounding errors in VPD because of the integration of multiple retrieved variables (Ts, Ta, U). FPAR was recovered with an RMS error of 2.4% over a wide range of sites in Oregon (Goward et al., 1994). There was some evidence of a lag between the CSI and soil moisture at depth at sites in Oregon. The CSI was found to predict surface soil moisture at a grassland site in central Kansas with an RMS error of 3.2% (Goetz, 1997).

10.2.4 Additional Quality Assessments

Although the results sometimes had low absolute accuracies, the field data themselves are not without error: although the inferences were usually for a >1 km² area made instantaneously, they were compared with point field values generally not measured at exactly the same times in the day or covering the same spatial area. Maps of retrieved variables had good relative accuracy and possibly better absolute accuracy than the comparisons with point measurements suggest.

10.2.5 Data Verification by Data Center

Data were examined for general consistency and clarity.

11. Notes

11.1 Limitations of the Data

The model is probably limited in terms of Ra recovery; hence NPP, because of the potential insensitivity of Ra to biomass in boreal forest stands (Goetz and Prince, 1998).

11.2 Known Problems with the Data

None.

11.3 Usage Guidance

Before uncompressing the Gzip files on CD-ROM, be sure that you have enough disk space to hold the uncompressed data files. Then use the appropriate decompression program provided on the CD-ROM for your specific system.

11.4 Other Relevant Information

Contact Dr. Scott J. Goetz for a platform-independent version of the model that is available; however, the model is highly system-tailored and requires 50 Gigabytes of space for a single run. If someone really wants it, Dr. Goetz could discuss with them some collaborative effort to get it functioning in another lab.

12. Application of the Data Set

The model may be operated with any remotely sensed observations that provide a measure of vegetation amount (e.g., spectral vegetation indices) and thermal emission in more than one channel (in order to get split-window surface radiometric temperature). The results are applicable to many studies, from characterizing carbon flux and storage over large areas to monitoring changes in forest productivity, stress, and management.

13. Future Modifications and Plans

A heterotrophic respiration (Rh) component is being added to the model in order to simulate, when combined with NPP, net ecosystem productivity (NEP). Simulated NEP is more directly comparable with eddy correlation (e.g., tower) carbon flux measurements than is NPP because of the lack of separability between the Ra and Rh components of measured soil CO₂ efflux. Moreover, NEP is important to characterize in order to quantify the direction of vegetation - atmosphere fluxes and to address carbon budgets at the local to global scale.

14. Software

14.1 Software Description

Version 1 of the model was developed and operated under the PCI image processing package using the Engineering Analysis and Scientific Interface (EASI) procedure language. The model has since been exported, primarily for speed of processing and modularity, to the UNIX environment. Version 2 is written in the C programming language. Version 2 is operable and changes to the model are reported in the upcoming publication Goetz et al. (1999a). BOREAS results with version 2 of the model are reported in Goetz et al. (1999b).

Gzip (GNU zip) uses the Lempel-Ziv algorithm (Welch, 1994) used in the zip and PKZIP commands.

14.2 Software Access

Contact Dr. Scott J. Goetz for a platform-independent version of the model that is available, however the model is highly system-tailored and requires 50 Gigabytes of space for a single run. If someone really wants it, Dr. Goetz could discuss with them some collaborative effort to get it functioning in another lab.

Gzip is available from many Web sites across the Internet (for example, ftp site prep.ai.mit.edu/pub/gnu/gzip-*.*) for a variety of operating systems in both executable and source code form. Versions of the decompression software for various systems are included on the CD-ROMs.

15. Data Access

The production efficiency model images are available from the Earth Observing System Data and Information System (EOSDIS) Oak Ridge National Laboratory (ORNL) Distributed Active Archive Center (DAAC).

15.1 Contact Information

For BOREAS data and documentation please contact:

ORNL DAAC User Services Oak Ridge National Laboratory P.O. Box 2008 MS-6407 Oak Ridge, TN 37831-6407 Phone: (423) 241-3952

Fax: (423) 574-4665

E-mail: ornldaac@ornl.gov or ornl@eos.nasa.gov

15.2 Data Center Identification

Earth Observing System Data and Information System (EOSDIS) Oak Ridge National Laboratory (ORNL) Distributed Active Archive Center (DAAC) for Biogeochemical Dynamics http://www-eosdis.ornl.gov/.

15.3 Procedures for Obtaining Data

Users may obtain data directly through the ORNL DAAC online search and order system [http://www-eosdis.ornl.gov/] and the anonymous FTP site [ftp://www-eosdis.ornl.gov/data/] or by contacting User Services by electronic mail, telephone, fax, letter, or personal visit using the contact information in Section 15.1.

15.4 Data Center Status/Plans

The ORNL DAAC is the primary source for BOREAS field measurement, image, GIS, and hardcopy data products. The BOREAS CD-ROM and data referenced or listed in inventories on the CD-ROM are available from the ORNL DAAC.

16. Output Products and Availability

16.1 Tape Products

None.

16.2 Film Products

None.

16.3 Other Products

These images are available on the BOREAS CD-ROM series.

Output image products listed in Section 7.4, Table 2, and described herein, that are not included on the BOREAS CD-ROM might be available on request.

A description of the model, summary of research papers, and results of model application at the global scale are available at the following URL: http://www.geog.umd.edu/glopem/.

17. References

17.1 Platform/Sensor/Instrument/Data Processing Documentation

Thawley, M. and J. Small. 1997. The Global Production Efficiency Model, version 2.0 (GLO-PEM2).

Welch, T.A. 1984. A Technique for High Performance Data Compression. IEEE Computer, Vol. 17, No. 6, pp. 8-19.

17.2 Journal Articles and Study Reports

Asrar, G., E.T. Kanemasu, R.D. Jackson, and P.J. Pinter. 1985. Estimation of total above ground phytomass production using remotely sensed data. Remote Sensing of Environment, 17: 211-220.

Becker, F. and Z.L. Li. 1990. Towards a local split window method over land surfaces. International Journal of Remote Sensing, 11: 1509-1522.

Collatz, G.J., J.T. Ball, C. Grivet, and J.A. Berry. 1991. Physiological and environmental regulation of stomatal conductance, photosynthesis and transpiration: a model that includes a laminar boundary layer. Agricultural and Forest Meteorology, 54: 107-136.

- Czajkowski, K.P., T. Mulhern, S.N. Goward, J. Cihlar, R.O. Dubayah, and S.D. Prince. 1997a. Biospheric environmental monitoring at BOREAS with AVHRR observations. Journal of Geophysical Research, 102 (D24): 29,651-29,662.
- Czajkowski, K.P., T. Mulhern, S.N. Goward, and J. Cihlar. 1997b. Validation of the geocoding and compositing system (GEOCOMP) using contextural analysis for AVHRR images. International Journal of Remote Sensing, 18 (14): 3055-3068.
- Goetz, S.J. 1997. Multi-sensor analysis of NDVI, surface temperature, and biophysical variables at a mixed grassland site. International Journal of Remote Sensing, 18: 71-94.
- Goetz, S.J. and S.D. Prince. 1996. Remote sensing of net primary production in boreal forest stands. Agricultural and Forest Meteorology, 78: 149-179.
- Goetz, S.J. and S.D. Prince. 1998. Variability in carbon exchange and light utilization among boreal forest stands: implications for remote sensing of net primary production. Canadian Journal of Forest Research 28: 375-389.
- Goetz, S.J. and S.D. Prince. 1999. Modeling terrestrial carbon exchange and storage: the evidence for and implications of functional convergence in light use efficiency. Advances in Ecological Research, 28: 57-92.
- Goetz, S.J., S.D. Prince, M.M. Thawley, K.P. Czajkowski, and S.N. Goward. 1999. Mapping regional patterns of canopy carbon exchange and storage with remote sensing. Journal of Geophysical Research (in press).
- Goetz, S.J., S.D. Prince, S.N. Goward, M. Thawley, and J. Small. 1999a. Satellite remote sensing of primary production: an improved production efficiency modeling approach. Ecological Modeling (in press).
- Goetz, S.J., S.D. Prince, S.N. Goward, M.M. Thawley, J. Small, and A. Johnston. 1999b. Mapping net primary production and related biophysical variables with remote sensing: application to the BOREAS region. Journal of Geophysical Research (in press).
- Goward, S.N. and D.G. Dye. 1997. Global biospheric monitoring with remote sensing. In: The use of remote sensing in modeling forest productivity (eds. H.L. Gholtz, K. Nakane, and H. Shimoda). pages 241-272, Kluwer Academic, New York.
- Goward, S.N., R.H. Waring, and D.G. Dye. 1994. Ecological remote sensing at OTTER: Macroscale satellite observations. Ecological Applications, 4: 322-343.
- Hall, F.G., Y. Shimabukuro, and K.F. Huemmrich. 1995. Remote sensing of forest biophysical structure in boreal stands of Picea mariana using mixture decomposition and geometric reflectance models. Ecological Applications, 5: 993-1013.
- Hanan, N.P., S.D. Prince, and A. Bèguè. 1995. Estimation of absorbed photosynthetically active radiation and vegetation net production efficiency using satellite data. Agricultural and Forest Meteorology, 76: 259-276.
- Hanan, N.P., S.D. Prince, and A. Bèguè. 1997. Modelling vegetation primary production during HAPEX-Sahel using production efficiency and canopy conductance model formulations. Journal of Hydrology, 188/189: 651-675.

- Kumar, M. and J.L. Monteith. 1982. Remote sensing of plant growth. In: Plants and the Daylight Spectrum (eds. H. Smith), pp. 133-144. Academic Press, London.
- Leemans, R. and W.P. Cramer. 1991. The IISAS database for mean monthly values of temperature, precipitation and cloudiness on a global terrestrial grid, Laxenburg, Austria International Institute for Applied Systems Analysis.
- Newcomer, J., D. Landis, S. Conrad, S. Curd, K. Huemmrich, D. Knapp, A. Morrell, J. Nickeson, A. Papagno, D. Rinker, R. Strub, T. Twine, F. Hall, and P. Sellers, eds. 2000. Collected Data of The Boreal Ecosystem-Atmosphere Study. NASA. CD-ROM.
- Potter, C.S., J.T. Randerson, C.B. Field, P.A. Matson, P.M. Vitousek, H.A. Mooney, and S.A. Klooster. 1993. Terrestrial ecosystem production: A process model based on global satellite and surface data. Global Biogeochemical Cycles, 7: 811-841.
- Prabhakara, C., G. Dalu, R.C. Lo, and R. Nath. 1977. Remote sensing of seasonal distribution of precipitable water over the oceans and inference of boundary layer structure. Monthly Weather Review, 107: 1388-1401.
- Prihodko, L. and S.N. Goward. 1997. Estimation of air temperature from remotely sensed observations. Remote Sensing of Environment, 60: 335-346.
- Prince, S.D. and S.J. Goward. 1995. Global primary production: a remote sensing approach. Journal of Biogeography, 22: 815-835.
- Prince, S.D., S.J. Goetz, and S.N. Goward. 1995. Monitoring primary production from Earth observing satellites. Water, Air, and Soil Pollution, 82: 509-522.
- Prince, S.D., S.J. Goetz, K. Czajkowski, R. Dubayah, and S.N. Goward. 1998. Inference of surface and air temperature, atmospheric precipitable water and vapor pressure deficit using AVHRR satellite observations: validation of algorithms. Journal of Hydrology, 213 (1-4), 230-249.
- Sellers, P. and F. Hall. 1994. Boreal Ecosystem-Atmosphere Study: Experiment Plan. Version 1994-3.0, NASA BOREAS Report (EXPLAN 94).
- Sellers, P. and F. Hall. 1996. Boreal Ecosystem-Atmosphere Study: Experiment Plan. Version 1996-2.0, NASA BOREAS Report (EXPLAN 96).
- Sellers, P., F. Hall, and K.F. Huemmrich. 1996. Boreal Ecosystem-Atmosphere Study: 1994 Operations. NASA BOREAS Report (OPS DOC 94).
- Sellers, P., F. Hall, and K.F. Huemmrich. 1997. Boreal Ecosystem-Atmosphere Study: 1996 Operations. NASA BOREAS Report (OPS DOC 96).
- Sellers, P., F. Hall, H. Margolis, B. Kelly, D. Baldocchi, G. den Hartog, J. Cihlar, M.G. Ryan, B. Goodison, P. Crill, K.J. Ranson, D. Lettenmaier, and D.E. Wickland. 1995. The boreal ecosystem-atmosphere study (BOREAS): an overview and early results from the 1994 field year. Bulletin of the American Meteorological Society. 76(9):1549-1577.
- Sellers, P.J., F.G. Hall, R.D. Kelly, A. Black, D. Baldocchi, J. Berry, M. Ryan, K.J. Ranson, P.M. Crill, D.P. Lettenmaier, H. Margolis, J. Cihlar, J. Newcomer, D. Fitzjarrald, P.G. Jarvis, S.T. Gower, D. Halliwell, D. Williams, B. Goodison, D.E. Wickland, and F.E. Guertin. 1997. BOREAS in 1997: Experiment Overview, Scientific Results and Future Directions. Journal of Geophysical Research 102(D24): 28,731-28,770.

Smith, W.L. 1966. Note on the relationship between precipitable water and surface dew point. Journal of Applied Meteorology, 5: 726-727.

Tucker, C.J., C.L. Vanpraet, E. Boerwinkel, and A. Gaston. 1983. Satellite remote sensing of total dry matter accumulation in the Senegalese Sahel. Remote Sensing of Environment, 13: 461-474.

17.3 Archive/DBMS Usage Documentation None.

18. Glossary of Terms

CSI - Cumulative surface wetness index

Gs - Stomatal conductance
e - Carbon yield of APAR
Ra - Autotrophic respiration
Rh - Heterotrophic respiration
Ta - Ambient air temperature

Ts - Surface radiometric temperature

U - Atmospheric precipitable water vapor amount

W - Standing above-ground biomass

19. List of Acronyms

APAR - Absorbed Photosynthetically Active Radiation

ASCII - American Standard Code for Information Interchange

AVHRR - Advanced Very High Resolution Radiometer Boreal-PEM - Boreal Forest Production Efficiency Model

BOREAS - BOReal Ecosystem-Atmosphere Study

BORIS - BOREAS Information System

CASA - Carnegie Ames Stanford Approach
CCRS - Canada Centre Remote Sensing
CD-ROM - Compact Disk-Read-Only memory
DAAC - Distributed Active Archive Center

EASI - Engineering Analysis and Scientific Interface

EOS - Earth Observing System

EOSDIS - EOS Data and Information System

FIFE - First International Satellite Land Surface Climatology Field

Experiment

FPAR - Fraction of incident PAR intercepted or absorbed

GEOCOMP - Geocoding and Compositioning System

GEWEX - Global Water Energy and Water Cycle Experiment

GIS - Geographic Information System
GLO-PEM - Global Production Efficiency Model

GOES - Geostationary Operational Environmental Satellite

GPP - Gross Primary Production

GSFC - Goddard Space Flight Center (NASA)

HAPEX-Sahel - Hydrology Atmosphere Pilot Experiment in the Sahel

HTML - HyperText Markup Language

IFC - Intensive Field Campaign

Lagal Area Coverage (of A)

LAC - Local Area Coverage (of AVHRR) NAD83 - North American Datum of 1983

NASA - National Aeronautics and Space Administration

NDVI - Normalized Difference Vegetation Index

NEP - Net Ecosystem Production

NOAA - National Oceanic and Atmospheric Administration

NPP - Net Primary Production NSA - Northern Study Area

ORNL - Oak Ridge National Laboratory
OTTER - Oregon Transect Experiment
PANP - Prince Albert National Park

PAR - Photosynthetically Active Radiation

PEM - Production Efficiency Model
RMS - Root Mean Square (Error)

SNF - Superior National Forest, Minnesota

SSA - Southern Study Area
TE - Terrestrial Ecology

TF - Tower Flux

URL - Uniform Resource Locator
UTM - Universal Transverse Mercator

VPD - Vapor Pressure Deficit

20. Document Information

20.1 Document Revision Date

Written: 25-Jun-1997

Last Updated: 24-Sep-1999

20.2 Document Review Date(s)

BORIS Review: 04-May-1999 Science Review: 26-Oct-1998

20.3 Document ID

20.4 Citation

When using these data, please include the following reference:

Goetz, S.J. and S.D. Prince. 1998. Variability in carbon exchange and light utilization among boreal forest stands: implications for remote sensing of net primary production. Canadian Journal of Forest Research 28: 375-389.

If using data from the BOREAS CD-ROM series, also reference the data as:

Goward, S.N., S.D. Prince, S.J. Goetz, K. Czajkowski, and R.O. Dubayah, "Biospheric Dynamics in the Boreal Forest Ecotone." In Collected Data of The Boreal Ecosystem-Atmosphere Study. Eds. J. Newcomer, D. Landis, S. Conrad, S. Curd, K. Huemmrich, D. Knapp, A. Morrell, J. Nickeson, A. Papagno, D. Rinker, R. Strub, T. Twine, F. Hall, and P. Sellers. CD-ROM. NASA, 2000.

Also, cite the BOREAS CD-ROM set as:

Newcomer, J., D. Landis, S. Conrad, S. Curd, K. Huemmrich, D. Knapp, A. Morrell, J. Nickeson, A. Papagno, D. Rinker, R. Strub, T. Twine, F. Hall, and P. Sellers, eds. Collected Data of The Boreal Ecosystem-Atmosphere Study. NASA. CD-ROM. NASA, 2000.

20.5 Document Curator

20.6 Document URL

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204. Adjington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

Davis Highway, Suite 1204, Arlington, VA 22202-430	22, and to the Office of Management and	Budget, Paperwork Reduction	Project (0704-0188), Washington, DC 20503.
1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE October 2000	3. REPORT TYPE AN Technical Me	
4. TITLE AND SUBTITLE	Getobel 2000	Technical Me	5. FUNDING NUMBERS
	amaal Eagsystam Atmassham	Study (DODEAC)	5. FUNDING NUMBERS
Technical Report Series on the Bo		-	
BOREAS TE-17 Production I	Efficiency Model Images		923
6. AUTHOR(S)			RTOP: 923-462-33-01
Scott J. Goetz, Samual N. Gov	ward, Stephen D. Prince,	Kevin Czajkowski.	
and Ralph O. Dubayah			
Forrest G. Hall and Andrea Pa	1 0 7		
7. PERFORMING ORGANIZATION NAME	E(S) AND ADDRESS (ES)		8. PEFORMING ORGANIZATION REPORT NUMBER
Goddard Space Flight Center			KEI OKI NOMBEK
Greenbelt, Maryland 20771			2000-03136-0
,			
9. SPONSORING / MONITORING AGE	NCY NAME(S) AND ADDRESS	S (ES)	10. SPONSORING / MONITORING
			AGENCY REPORT NUMBER
National Aeronautics and Space	Administration		TM-2000-209891
Washington, DC 20546-0001			Vol. 171
11. SUPPLEMENTARY NOTES			
S. Goetz, S.N. Goward, S.D.	Prince K Czaikoweki a	nd P O Dubayah	University of Maryland
	, ,	•	t Center, Greenbelt, Maryland
Conege Park, A. Papagno. Ka	iyulcoli 1155, NASA Go	ddaid Space Fiigii	t Center, Greenbert, Waryland
12a. DISTRIBUTION / AVAILABILITY STA	TEMENT		12b. DISTRIBUTION CODE
Unclassified-Unlimited			
Subject Category: 43			
Report available from the NASA	A Center for AeroSpace Inf	formation,	
7121 Standard Drive, Hanover, 1	MD 21076-1320. (301) 62	1-0390.	
13. ABSTRACT (Maximum 200 words)			
	obal Production Efficien	cv Model (http://w	ww.inform.umd.edu/glopem/

A BOREAS version of the Global Production Efficiency Model (http://www.inform.umd.edu/glopem. [Internet Link]) was developed by TE-17 to generate maps of gross and net primary production, autotrophic respiration, and light use efficiency for the BOREAS region. This document provides basic information on the model and how the maps were generated. The data generated by the model are stored in binary image-format files.

14. SUBJECT TERMS BOREAS, terrestrial eco	logy.		15. NUMBER OF PAGES 19
			16. PRICE CODE
17. SECURITY CLASSIFICATION OF REPORT	18. SECURITY CLASSIFICATION OF THIS PAGE	19. SECURITY CLASSIFICATION OF ABSTRACT	20. LIMITATION OF ABSTRACT
Unclassified	Unclassified	Unclassified	UL